

LECTURES AT UIC

WENLIANG ZHANG

Throughout this lecture, p denotes a prime number and R denotes a commutative noetherian ring of characteristic p .

Definition 0.1. The *Frobenius endomorphism* of R is the map

$$f : R \xrightarrow{r \mapsto r^p} R.$$

Since R has characteristic p , it follows from the binomial formula that $(a+b)^p = a^p + b^p$ and consequently f is a ring homomorphism, and so are its iterates f^e for each integer $e \geq 1$.

We will use $F_*^e R$ to denote the R -module that is the same as R as an abelian group and whose R -module structure is defined via $F^e : R \rightarrow R$. The Peskine-Szpiro functor, denoted by F_R , is defined via

$$F_R(M) := F_* R \otimes_R M.$$

The e -th iteration of F_R is denoted by F_R^e , which is clearly given by $F_R^e(M) = F_*^e R \otimes_R M$.

Exercise 0.2. Prove that

- (1) $F_R^e(R^\ell) \cong R^\ell$ for all free modules R^ℓ ;
- (2) $F_R^e(R/I) \cong R/I^{[p^e]}$ for each ideal I of R ;
- (3) $F_R^e(R_g) \cong R_g$ for each element $g \in R$;
- (4) F_R commutes with localization, i.e., $F_R(M)_p \cong F_{R_p}(M_p)$.

Let $\mathfrak{a} = (g_1, \dots, g_n)$ be an ideal of R and $\varphi : R \rightarrow S$ be a ring homomorphism between commutative noetherian rings of characteristic p . Tensoring φ with the Čech complex $\check{C}(R; \underline{g})$ produces a map of complexes:

$$\check{C}(R; \underline{g}) \rightarrow \check{C}(S; \varphi(\underline{g})).$$

Therefore, there is a homomorphism of R -modules

$$H_{\mathfrak{a}}^i(R) \rightarrow H_{\varphi(\mathfrak{a})S}^i(S).$$

Specializing in the case when $S = R$ and $\varphi = f$, we have the *Frobenius action* on $H_{\mathfrak{a}}^i(R)$:

$$H_{\mathfrak{a}}^i(R) \rightarrow H_{f(\mathfrak{a})R}^i(R) = H_{\mathfrak{a}}^i(R),$$

where the equality follows from $f(\mathfrak{a}) = (g_1^p, \dots, g_n^p)$ which has the same radical as \mathfrak{a} . We will denote this Frobenius action by F .

Example 0.3. Let $R = k[x_1, \dots, x_d]$ (or $R = k[[x_1, \dots, x_d]]$) and $\mathfrak{a} = (x_1, \dots, x_d)$ where k is a field of characteristic p . The Frobenius action on $H_{\mathfrak{a}}^d(R)$ is given by

$$\left[\frac{c}{x_1^{e_1} \dots x_d^{e_d}} \right] \mapsto \left[\frac{c^p}{x_1^{pe_1} \dots x_d^{pe_d}} \right],$$

where $c \in k$.

Example 0.4. Let (R, \mathfrak{m}) be a noetherian local ring and x_1, \dots, x_d be a system of parameters, where $d = \dim(R)$. Then each element of $H_{\mathfrak{m}}^d(R)$ can be written as $[\frac{r}{x_1^{e_1} \dots x_d^{e_d}}]$ for some $r \in R$ and positive integers e_i . The Frobenius action on $H_{\mathfrak{m}}^d(R)$ is given by

$$[\frac{r}{x_1^{e_1} \dots x_d^{e_d}}] \mapsto [\frac{r^p}{x_1^{pe_1} \dots x_d^{pe_d}}].$$

One may notice that, in the case when $R = k[x_1, \dots, x_d]$, one has R is a free module over $f(R) = k^p[x_1^p, \dots, x_d^p]$; hence $f : R \rightarrow R$ is flat. More generally, we have the following theorem due to Kunz [Kun69].

Theorem 0.5 (Kunz). *The following are equivalent:*

- (1) R is regular;
- (2) f^e is a flat ring homomorphism for each $e \geq 1$;
- (3) f^e is a flat ring homomorphism for some $e \geq 1$.

Theorem 0.6. *Assume that R is regular and \mathfrak{a} is an ideal of R . Then*

$$F_R^e(H_{\mathfrak{a}}^i(R)) \cong H_{\mathfrak{a}}^i(R).$$

Proof. Since $H_{\mathfrak{a}}^i(R) = \varinjlim_n \text{Ext}^i(R/\mathfrak{a}^n, R)$, we have

$$\begin{aligned} F_*^e R \otimes_R H_{\mathfrak{a}}^i(R) &= F_*^e R \otimes_R \varinjlim_n \text{Ext}^i(R/\mathfrak{a}^n, R) \\ &\cong \varinjlim_n F_*^e R \otimes_R \text{Ext}^i(R/\mathfrak{a}^n, R) \\ &\cong \varinjlim_n \text{Ext}^i(F_*^e R \otimes_R R/\mathfrak{a}^n, F_*^e R \otimes_R R) \text{ since } F_*^e R \text{ is flat} \\ &\cong \varinjlim_n \text{Ext}^i(R/(\mathfrak{a}^{[p^e]})^n, R) \text{ by Exercise 0.2} \\ &\cong H_{\mathfrak{a}^{[p^e]}}^i(R) \\ &= H_{\mathfrak{a}}^i(R) \end{aligned}$$

Alternatively, one may also prove it via the Čech complex characterization of $H_{\mathfrak{a}}^i(R)$, using the facts that $F_*^e R$ is flat and that $F_R^e(R_{\mathfrak{g}}) \cong R_{\mathfrak{g}}$. \square

Exercise 0.7. Prove that, if R is regular, then $F_R(E) \cong E$ for each injective R -module E .

1. VANISHING OF LOCAL COHOMOLOGY MODULES

Exercise 1.1. Let (R, \mathfrak{m}) be a regular local ring of characteristic p . Assume that $\text{Ann}_R(H_{\mathfrak{a}}^i(R)) \neq 0$. Prove $H_{\mathfrak{a}}^i(R) = 0$.

Exercise 1.2. Let \mathfrak{p} be a prime ideal of a regular ring R of characteristic p . Prove that $\mathfrak{p}^{[p^e]}$ is \mathfrak{p} -primary.

Theorem 1.3 (Peskin-Szpiro [PS73]). *Let (R, \mathfrak{m}) be a regular local ring of characteristic p of dimension n and let I be an ideal of R . Assume that $\text{depth}(R/I) \geq c$. Then*

$$H_I^{n-i}(R) = 0 \text{ for } i \leq c - 1.$$

Proof. Since $\text{depth}(R/I) \geq c$, we have $H_{\mathfrak{m}}^i(R/I) = 0$ for $i \leq c - 1$. By local duality, $\text{Ext}_R^{n-i}(R/I, R) = 0$. By Theorem 0.5, we have $\text{Ext}_R^{n-i}(R/I^{[p^e]}, R) \cong F_*^e R \otimes_R \text{Ext}_R^{n-i}(R/I, R) = 0$. Hence

$$H_I^{n-i}(R) = \varinjlim_e \text{Ext}_R^{n-i}(R/I^{[p^e]}, R) = 0.$$

□

Example 1.4. Let $R = k[[x_{ij}]]$ with $i = 1, 2$ and $j = 1, 2, 3$ where k is a field. Let I be generated by the 2×2 minors of (x_{ij}) . Then R/I has depth 4 by [HE71].

When k has characteristic p , by Theorem 1.3, we have $H_I^i(R) \neq 0$ if and only if $i = 2$.

In contrast, when k has characteristic 0, one has $H_I^3(R) \neq 0$ ([HS77, page 75]).

Exercise 1.5. Let (R, \mathfrak{m}) be a complete regular local ring and E be the injective hull of the residue field R/\mathfrak{m} . Denote the Matlis dual functor $\text{Hom}_R(-, E)$ by $D(-)$. Prove that there is a functorial R -module isomorphism

$$\tau : D(F_R(M)) \cong F_R(D(M))$$

for all artinian R -modules M .

Let A be a noetherian commutative ring of characteristic p . Let $A\{f\}$ be the subring of $\text{Hom}_{\mathbb{Z}}(A, A)$ generated by multiplications by elements of A and $f : A \xrightarrow{a \mapsto a^p} A$, or equivalently $A\{f\} = \frac{A\langle f \rangle}{(fa - a^p f)}$. An $A\{f\}$ -module M is an A -module M equipped with a \mathbb{Z} -linear map $f : M \rightarrow M$ such that $f(am) = a^p f(m)$ for all $a \in A$ and $m \in M$.

Let (R, \mathfrak{m}) be a complete regular local ring and A be a homomorphic image of R . Let M be an $A\{f\}$ -module. One can check that

$$\alpha : F_R(M) \xrightarrow{r \otimes m \mapsto r f(m)} M$$

is an R -module homomorphism. Now, assume that M is artinian. Taking the Matlis dual of α and applying Exercise 1.5, we have an R -module homomorphism

$$\beta = \tau \circ D(\alpha) : D(M) \rightarrow F_R(D(M)),$$

and hence we have a direct system of R -modules:

$$D(M) \xrightarrow{\beta} F_R(D(M)) \xrightarrow{F_R(\beta)} F_R^2(D(M)) \rightarrow \dots$$

Definition 1.6 (Lyubeznik's $\mathcal{H}_{R,A}$ functor [Lyu97]). Let (R, \mathfrak{m}) be a complete regular local ring and A be a homomorphic image of R . For each artinian $A\{f\}$ -module, we define

$$\mathcal{H}_{R,A}(M) := \varinjlim (D(M) \xrightarrow{\beta} F_R(D(M)) \xrightarrow{F_R(\beta)} F_R^2(D(M)) \rightarrow \dots)$$

Example 1.7. Let (R, \mathfrak{m}) be a complete regular local ring of dimension n and A be a homomorphic image of R . Let I be the kernel of $R \twoheadrightarrow A$. From our discussion, we know that $H_{\mathfrak{m}}^i(A)$ is naturally an $A\{f\}$ -module. We would like to understand $\mathcal{H}_{R,A}(H_{\mathfrak{m}}^i(A))$.

By local duality, we have $D(H_{\mathfrak{m}}^i(A)) \cong \text{Ext}_R^{n-i}(R/I, R)$ and hence the direct system

$$\text{Ext}_R^{n-i}(R/I, R) \xrightarrow{\beta} F_R(\text{Ext}_R^{n-i}(R/I, R)) \xrightarrow{F_R(\beta)} F_R^2(\text{Ext}_R^{n-i}(R/I, R)) \rightarrow \dots$$

which is the same as

$$\text{Ext}_R^{n-i}(R/I, R) \rightarrow \text{Ext}_R^{n-i}(R/I^{[p]}, R) \rightarrow \text{Ext}_R^{n-i}(R/I^{[p^2]}, R) \rightarrow \dots$$

whose direct limit is $H_I^{n-i}(R)$. Therefore,

$$\mathcal{H}_{R,A}(H_m^i(A)) \cong H_I^{n-i}(R).$$

Theorem 1.8 (Theorem 1.1 in [Lyu06]). *Let (R, \mathfrak{m}) be a complete regular local ring of dimension n and A be a homomorphic image of R . Let I be the kernel of $R \rightarrow A$. Then $H_I^{n-i}(R) = 0$ if and only if the Frobenius action on $H_m^i(A)$ is nilpotent.*

Proof. First note that, since

$$H_i^{n-i}(R) = \varinjlim_e \text{Ext}_R^{n-i}(R/I^{[p^e]}, R)$$

and

$$F_R^e(\text{Ext}_R^{n-i}(R/I, R)) \cong \text{Ext}_R^{n-i}(R/I^{[p^e]}, R),$$

we see that $H_i^{n-i}(R) = 0$ if and only if there is t such that

$$\varphi_t : \text{Ext}_R^{n-i}(R/I, R) \rightarrow \text{Ext}_R^{n-i}(R/I^{[p^t]}, R)$$

is the 0 map.

With notation as in Example 1.7, $\varphi_t = 0$ if and only if $F_R^{t-1}(\beta) \circ \cdots \circ F_R(\beta) \circ \beta = 0$. Since $\text{Ext}_R^{n-i}(R/I, R)$ is finitely generated (or equivalently $H_m^i(R/I)$ is artinian), by Matlis duality, $F_R^{t-1}(\beta) \circ \cdots \circ F_R(\beta) \circ \beta = 0$ if and only if the following composition of maps is 0

$$F_R^t(H_m^i(R/I)) \xrightarrow{F_R^{t-1}(\alpha)} F_R^{t-1}(H_m^i(R/I)) \rightarrow \cdots \rightarrow F_R(H_m^i(R/I)) \xrightarrow{\alpha} H_m^i(R/I).$$

Writing $F_R^t(H_m^i(R/I))$ as $F_*R \otimes_R \cdots \otimes_R F_*R \otimes_R H_m^i(R/I)$, one can check that

$$\alpha \circ \cdots \circ F_R^{t-1}(\alpha)(r_t \otimes \cdots \otimes r_1 \otimes z) = r_t r_{t-1}^p \cdots r_1^{p^t} f^t(z)$$

for all $r_i \in F_*R$ and $z \in H_m^i(R/I)$. It is now straightforward to check that $\alpha \circ \cdots \circ F_R^{t-1}(\alpha) = 0$ if and only if $f^t = 0$, *i.e.* the Frobenius action $f : H_m^i(R/I) \rightarrow H_m^i(R/I)$ is nilpotent. This completes the proof. \square

In light of Theorem 1.8, we introduce the notion of F -depth:

Definition 1.9. Let (A, \mathfrak{m}) be a noetherian local ring of characteristic p . The F -depth of A is the smallest integer j such that the Frobenius on $H_m^j(A)$ is nilpotent.

Exercise 1.10. Prove that $F\text{-depth}(A) \leq \dim(A)$.

As an application of Theorem 1.8, we will state (without proofs) a result on cohomological dimension of an open subset of \mathbb{P}_k^n , where k is a field of characteristic p . To this end, let's recall the definition of cohomological dimension.

Definition 1.11. Let $Y \subset \mathbb{P}_k^n$ be a closed subscheme and U be the complement of Y in \mathbb{P}_k^n . The cohomological dimension of U , denoted by $\text{cd}(U)$, is the largest integer i such there is a quasi-coherent sheaf \mathcal{F} on U such that $H^i(U, \mathcal{F}) \neq 0$.

Remark 1.12. Set $R = k[x_0, \dots, x_n]$ and I to be the homogeneous defining ideal of Y . Then it is proved in [Har68, pp. 412-413] that $\text{cd}(U) < c$ if and only if $H_I^j(R) = 0$ for all $j > c$.

Theorem 1.13 (Corollary 5.4 in [Lyu06]). *Let Y, U be the same as above. For each $c \geq 2$, we have $\text{cd}(U) < n - c$ if and only if the following 3 conditions hold:*

- (1) Y is geometrically connected and $\dim(Y) > 0$;
- (2) F -depth($\mathcal{O}_{Y,y}$) $\geq c$ for every closed point $y \in Y$;
- (3) the Frobenius action on $H^i(Y, \mathcal{O}_Y)$ is nilpotent for each $1 \leq j \leq c - 1$.

2. ASSOCIATED PRIMES AND BASS NUMBERS OF LOCAL COHOMOLOGY MODULES

A question of Huneke [Hun92, Problem 4] asks whether local cohomology modules of Noetherian rings have finitely many associated prime ideals. The answer is negative in general; the first counterexample was given by Singh [Sin00, § 4]. There are positive answers in some cases; for instance, when the ring is a regular ring of characteristic p . We want to discuss the approach in [HS93].

Recall that, when A is a commutative noetherian local ring and M is an A -module, the Bass number $\mu^i(\mathfrak{p}, M)$ (with respect to a prime ideal \mathfrak{p} of A) is defined as

$$\mu^i(\mathfrak{p}, M) := \dim_{\kappa(\mathfrak{p})}(\mathrm{Ext}_{A_{\mathfrak{p}}}^i(\kappa(\mathfrak{p}), M_{\mathfrak{p}})),$$

where $\kappa(\mathfrak{p}) = A_{\mathfrak{p}}/\mathfrak{p}A_{\mathfrak{p}}$. Alternatively, let $E^i(M)$ denote the i -th item in the minimal injective resolution of M . One has a decomposition $E^i(M) \cong \bigoplus_{\lambda \in \Lambda} E(A/\mathfrak{p}_{\lambda})$. Then for each prime \mathfrak{p} , one has $\mu^i(\mathfrak{p}, M)$ is the cardinality of the set $\{\lambda \in \Lambda \mid \mathfrak{p}_{\lambda} = \mathfrak{p}\}$.

Exercise 2.1. Prove that $\mathfrak{p} \in \mathrm{Ass}_A(M)$ if and only if $\mu^0(\mathfrak{p}, M) > 0$.

Exercise 2.2. Let (R, \mathfrak{m}) be a noetherian local ring and let $\{\mathcal{I}_j\}_{j \in \mathbb{Z}_{>0}}$ be a direct system of injective R -modules with limit \mathcal{I} . Assume that there is an integer μ such that each \mathcal{I}_j has at most μ copies of $E(R/\mathfrak{m})$. Prove \mathcal{I} also has at most μ copies of $E(R/\mathfrak{m})$.

Theorem 2.3 (Huneke-Sharp). *Let (R, \mathfrak{m}) be a regular local ring of characteristic p and I be an ideal of R . Then $\mu^i(\mathfrak{p}, H_I^j(R)) \leq \mu^i(\mathfrak{p}, \mathrm{Ext}_R^j(R/I, R))$ for each prime ideal \mathfrak{p} and all integers i, j . In particular, $\mu^i(\mathfrak{p}, H_I^j(R))$ is finite.*

Proof. After localizing at \mathfrak{p} , we may and we do assume that $\mathfrak{p} = \mathfrak{m}$. Let

$$0 \rightarrow \mathrm{Ext}_R^j(R/I, R) \rightarrow E^0 \rightarrow E^1 \rightarrow \dots$$

be a minimal injective resolution of $\mathrm{Ext}_R^j(R/I, R)$. For each positive integer t , we will denote $F_R^t(E^j)$ by E_t^j . By Exercise 0.7, we know that $E_t^j \cong E^j$ for each t . Applying the exact functor F_R^t to this injective resolution of $\mathrm{Ext}_R^j(R/I, R)$, we get an exact sequence

$$0 \rightarrow \mathrm{Ext}_R^j(R/I^{[p^t]}, R) \rightarrow E_t^0 \rightarrow E_t^1 \rightarrow \dots$$

which is a minimal injective resolution of $\text{Ext}_R^j(R/I^{[p^t]}, R)$. The functoriality of F_R and standard results on injective resolutions produce a commutative diagram

$$\begin{array}{ccccccc}
0 & \longrightarrow & \text{Ext}_R^j(R/I, R) & \longrightarrow & E^0 & \longrightarrow & \cdots \longrightarrow E^j \longrightarrow \cdots \\
& & \downarrow & & \downarrow & & \downarrow \\
0 & \longrightarrow & \text{Ext}_R^j(R/I^{[p]}, R) & \longrightarrow & E_1^0 & \longrightarrow & \cdots \longrightarrow E_1^j \longrightarrow \cdots \\
& & \downarrow & & \downarrow & & \downarrow \\
& & \vdots & & \vdots & & \vdots \\
& & \downarrow & & \downarrow & & \downarrow \\
0 & \longrightarrow & \text{Ext}_R^j(R/I^{[p^t]}, R) & \longrightarrow & E_t^0 & \longrightarrow & \cdots \longrightarrow E_t^j \longrightarrow \cdots \\
& & \downarrow & & \downarrow & & \downarrow \\
& & \vdots & & \vdots & & \vdots
\end{array}$$

Taking direct limits and keeping in mind that a direct of injective modules is still injective, we get an exact sequence

$$0 \rightarrow H_I^j(R) \rightarrow \mathcal{E}^0 \rightarrow \mathcal{E}^1 \rightarrow \cdots$$

which is a minimal injective resolution of $H_I^j(R)$. It follows from Exercise 2.2 and $E_t^j \cong E^j$ that

$$\mu^i(\mathfrak{p}, H_I^j(R)) \leq \mu^i(\mathfrak{p}, \text{Ext}_R^j(R/I, R)).$$

□

As an immediate consequence of Theorem 2.3 and Exercise 2.1, we have:

Theorem 2.4 (Huneke-Sharp). *Let R be a regular ring of characteristic p and I be an ideal of R . Then $\text{Ass}_R(H_I^j(R)) \subseteq \text{Ass}_R(\text{Ext}_R^j(R/I, R))$. In particular, $\text{Ass}_R(H_I^j(R))$ is a finite set.*

There are still many interesting question regarding ‘finiteness’ properties of local cohomology modules in characteristic p when R is not regular. Before we proceed to discussions of non-regular rings, I’d like to discuss Lyubeznik’s F -module theory, which is a generalization of what we have seen so far.

3. F -MODULES

Definition 3.1. Let R be a regular ring of characteristic p . An F -module is an R -module M equipped with an R -linear isomorphism $\theta_M : M \rightarrow F_R(M)$.

A homomorphism between F -modules (M, θ_M) and (N, θ_N) is a homomorphism $\varphi : M \rightarrow N$ such that the following is a commutative diagram

$$\begin{array}{ccc}
M & \xrightarrow{\theta_M} & F_R(M) \\
\downarrow \varphi & & \downarrow F_R(\varphi) \\
N & \xrightarrow{\theta_N} & F_R(N).
\end{array}$$

Example 3.2. It follows from Exercise 0.2 and Theorem 0.6 that R^ℓ , R_g , and $H_a^j(R)$ are F -modules.

From what we have seen, the morphism $\text{Ext}_R^j(R/\mathfrak{a}, R) \rightarrow F_R(\text{Ext}_R^j(R/\mathfrak{a}, R))$ plays a key role in understanding $H_a^j(R)$. The analog of such a morphism for an F -module is called a generating morphism.

Definition 3.3. Let R be a regular ring of characteristic p and \mathcal{M} be an F -module. A *generating morphism* of \mathcal{M} is an R -module homomorphism $M \rightarrow F_R(M)$, with some R -module M , such that \mathcal{M} is the direct limit of the top row of the following commutative diagram:

$$\begin{array}{ccccccc} M & \xrightarrow{\beta} & F_R(M) & \xrightarrow{F_R(\beta)} & F_R^2(M) & \xrightarrow{F_R^2(\beta)} & \dots \\ \downarrow \beta & & \downarrow F_R(\beta) & & \downarrow F_R^2(\beta) & & \\ F_R(M) & \xrightarrow{F_R(\beta)} & F_R^2(M) & \xrightarrow{F_R^2(\beta)} & F_R^3(M) & \xrightarrow{F_R^3(\beta)} & \dots \end{array}$$

and the structure isomorphism $\theta : \mathcal{M} \rightarrow F_R(\mathcal{M})$ is induced from the vertical arrows in the diagram.

Remark 3.4. Since tensor product commutes with direct limits, so the direct limit of the bottom row is indeed $F_R(\mathcal{M})$.

It is clear that each F -module \mathcal{M} has a generating morphism, the structure isomorphism.

Example 3.5. The morphism $\text{Ext}_R^j(R/\mathfrak{a}, R) \rightarrow F_R(\text{Ext}_R^j(R/\mathfrak{a}, R))$ is clearly a generating morphism for the F -module $H_a^j(R)$.

Given $f \in R$, there is a map of Koszul complexes

$$\begin{array}{ccccccc} K^\bullet(f; R) & = & 0 & \longrightarrow & R & \xrightarrow{f} & R & \longrightarrow & 0 \\ \downarrow & & & & \parallel & & \downarrow f^{p-1} & & \\ K^\bullet(f^p; R) & = & 0 & \longrightarrow & R & \xrightarrow{f^p} & R & \longrightarrow & 0 \end{array}$$

where K^\bullet denotes the Koszul complex. Let $\mathbf{f} = f_1, \dots, f_n$ be a sequence of elements of R . Regarding $K^\bullet(\mathbf{f}; R)$ as the tensor products

$$K^\bullet(f_1; R) \otimes \dots \otimes K^\bullet(f_n; R),$$

one obtains a map of complexes

$$K^\bullet(\mathbf{f}; R) \longrightarrow K^\bullet(\mathbf{f}^p; R),$$

and induced maps on cohomology modules

$$H^j(\mathbf{f}; R) \xrightarrow{\beta} H^j(\mathbf{f}^p; R),$$

where \mathfrak{a} is the ideal generated by \mathbf{f} . Since $H_a^j(R)$ is the direct limit of $\{H^j(\mathbf{f}^{p^t}; R)\}_t$ and $H^j(\mathbf{f}; R) \cong F_R(H^j(\mathbf{f}^p; R))$, the composition $H^j(\mathbf{f}; R) \rightarrow H^j(\mathbf{f}^p; R) \cong F_R(H^j(\mathbf{f}^p; R))$ is also a generating morphism of $H_a^j(R)$.

Note that $H_a^j(R)$ is ‘generated’ by a finitely generated R -module $\text{Ext}_R^j(R/\mathfrak{a}, R)$ (or $H^j(\mathbf{f}; R)$) via the generating morphism $\text{Ext}_R^j(R/\mathfrak{a}, R) \rightarrow F_R(\text{Ext}_R^j(R/\mathfrak{a}, R))$ (or $H^j(\mathbf{f}; R) \rightarrow H^j(\mathbf{f}^p; R)$). This leads to: ■

Definition 3.6. An F -module \mathcal{M} is called F -finite if \mathcal{M} admits a generating morphism $\beta : M \rightarrow F_R(M)$ with M a finitely generated R -module.

If, in addition, β is injective, M (or $\beta(M)$) is called a *root* of \mathcal{M} and β is called a *root morphism*.

Theorem 3.7. *Every F -finite module has a root.*

Proof. Let \mathcal{M} be an F -finite module with a generating morphism $\beta : M \rightarrow F_R(M)$ with M a finitely generated R -module. Let $\beta_i : M \rightarrow F_R^i(M)$ denote the composition $M \xrightarrow{\beta} F_R(M) \xrightarrow{F_R(\beta)} \dots \xrightarrow{F_R^{i-1}(\beta)} F_R^i(M)$. It is clear that $\ker(\beta_i) \subseteq \ker(\beta_{i+1})$ and hence $\ker(\beta_i) = \ker(\beta_{i+1}) = \dots$ for $i \gg 0$.

We claim that $\text{Im}(\beta_i)$ is a root of \mathcal{M} .

Let f denote the composition of $\text{Im}(\beta_i) \xrightarrow{\sim} \text{Im}(\beta_{i+1})$ and $\text{Im}(\beta_{i+1}) \hookrightarrow F_R(\text{Im}(\beta_i))$. Since F_R is exact, we have $\text{Im}(F_R^t(\beta_i)) = F_R^t(\text{Im}(\beta_i))$. Hence the direct system $M \xrightarrow{\beta} F_R(M) \xrightarrow{F_R(\beta)} \dots$ induces a direct system

$$\text{Im}(\beta_i) \xrightarrow{f} F_R(\text{Im}(\beta_i)) \xrightarrow{F_R(f)} F_R^2(\text{Im}(\beta_i)) \xrightarrow{F_R^2(f)} \dots$$

Then we have a commutative diagram of direct systems

$$\begin{array}{ccccccc} M & \xrightarrow{\beta} & F_R(M) & \xrightarrow{F_R(\beta)} & F_R^2(M) & \xrightarrow{F_R^2(\beta)} & \dots \\ \downarrow \beta_i & & \downarrow F_R(\beta_i) & & \downarrow F_R^2(\beta_i) & & \\ \text{Im}(\beta_i) & \xrightarrow{f} & F_R(\text{Im}(\beta_i)) & \xrightarrow{F_R(f)} & F_R^2(\text{Im}(\beta_i)) & \xrightarrow{F_R^2(f)} & \dots \end{array}$$

Since vertical maps are surjective, the induced map on the direct limits is surjective. The map on the direct limits is also injective since the kernel of each vertical map eventually goes to 0 in the direct system. Hence the direct limit of the bottom system is also \mathcal{M} . Since $\text{Im}(\beta_i)$ is a finite R -module and f is injective, $\text{Im}(\beta_i)$ is a root of \mathcal{M} . \square

Example 3.8. $H_{\mathfrak{a}}^j(R)$ has a root since it is an F -finite module.

One desirable feature of F -module is the following.

Theorem 3.9 (Theorem 1.4 in [Lyu97]). $\text{inj. dim}_R(\mathcal{M}) \leq \dim_R(\text{Supp}_R(\mathcal{M}))$ for each F -module \mathcal{M} .

Remark 3.10. An immediate consequence of Theorem 3.9 is that, if \mathfrak{m} is a maximal ideal of R , then $H_{\mathfrak{m}}^i(H_{\mathfrak{a}}^j(R))$ (or more generally, $H_{\mathfrak{m}}^i H_{\mathfrak{a}_1}^{j_1} \dots H_{\mathfrak{a}_s}^{j_s}(R)$) is a direct sum of some copies of $E(R/\mathfrak{m})$. This leaves the following question open: let $R = k[x_1, \dots, x_n]$ be the polynomial ring over a field k with the usual grading (*i.e.* $\deg(x_i) = 1$) and $\mathfrak{a}_1, \dots, \mathfrak{a}_s$ be graded ideals of R . Set $\mathfrak{m} = (x_1, \dots, x_n)$. Is it true that $H_{\mathfrak{m}}^i H_{\mathfrak{a}_1}^{j_1} \dots H_{\mathfrak{a}_s}^{j_s}(R) \cong E(R/\mathfrak{m})^{\oplus t}$ for some integer t as graded R -modules? Or, equivalently, are all the socle elements of $H_{\mathfrak{m}}^i H_{\mathfrak{a}_1}^{j_1} \dots H_{\mathfrak{a}_s}^{j_s}(R)$ have degree $-n$? We will come back to this question when we discuss \mathcal{D} -modules.

The following is a generalization of Theorem 2.3.

Theorem 3.11 (Theorem 2.11 in [Lyu97]). *Let \mathcal{M} be an F_R -finite module and M be a root of \mathcal{M} . Then*

$$\mu^j(\mathfrak{p}, \mathcal{M}) \leq \mu^j(\mathfrak{p}, M)$$

for each prime ideal \mathfrak{p} and each integer j .

Exercise 3.12. Let \mathfrak{p} be a prime ideal of R . Assume that M is an R -module such that $H_{\mathfrak{p}}^j(M)_{\mathfrak{p}}$ are injective for all j . Prove that

$$\mu^j(\mathfrak{p}, M) = \mu^0(\mathfrak{p}, H_{\mathfrak{p}}^j(M)).$$

Exercise 3.13. Prove Theorem 3.11.

Surprisingly, local cohomology modules, though may not be finitely generated or artinian as R -modules, will have finite length in the category of F_R -modules.

Theorem 3.14. *If R is a noetherian regular ring essentially of finite type over a regular local ring of characteristic p , then every F_R -finite module has finite length in the category of F_R -modules.*

4. \mathcal{D} -MODULES

Let R be a commutative ring. \mathbb{Z} -linear differential operators on R are defined inductively as follows: for each $r \in R$, the multiplication by r map $\tilde{r}: R \rightarrow R$ is a differential operator of order 0; for each positive integer n , the differential operators of order less than or equal to n are those additive maps $\delta: R \rightarrow R$ for which the commutator

$$[\tilde{r}, \delta] = \tilde{r} \circ \delta - \delta \circ \tilde{r}$$

is a differential operator of order less than or equal to $n - 1$. If δ and δ' are differential operators of order at most m and n respectively, then $\delta \circ \delta'$ is a differential operator of order at most $m + n$. Thus, the differential operators on R form a subring $\mathcal{D}(R)$ of $\text{End}_{\mathbb{Z}}(R)$.

When R is an algebra over a commutative ring A , we define $\mathcal{D}(R, A)$ to be the subring of $\mathcal{D}(R)$ consisting of differential operators that are A -linear. Note that $\mathcal{D}(R, \mathbb{Z}) = \mathcal{D}(R)$.

Example 4.1. Let $R = A[x_1, \dots, x_d]$ or $A[[x_1, \dots, x_d]]$. Then $\frac{1}{t_i!} \frac{\partial^{t_i}}{\partial x_i^{t_i}}$ can be viewed as a differential operator on R even if the integer $t_i!$ is not invertible. In either case, $\mathcal{D}(R, A)$ is the free R -module with basis

$$\frac{1}{t_1!} \frac{\partial^{t_1}}{\partial x_1^{t_1}} \cdots \frac{1}{t_d!} \frac{\partial^{t_d}}{\partial x_d^{t_d}} \quad \text{for } (t_1, \dots, t_d) \in \mathbb{N}^d,$$

or equivalently $\mathcal{D}(R, A) = R \langle \frac{1}{t_1!} \frac{\partial^{t_1}}{\partial x_1^{t_1}} \cdots \frac{1}{t_d!} \frac{\partial^{t_d}}{\partial x_d^{t_d}} \mid (t_1, \dots, t_d) \in \mathbb{N}^d \rangle$

By the quotient rule, it is clear that R_f is a $\mathcal{D}(R, A)$ -module. It is also straightforward to check that the maps in the Čech complex are $\mathcal{D}(R, A)$ -linear and hence each local cohomology module $H_{\mathfrak{a}}^i(R)$ is a $\mathcal{D}(R, A)$ -module.

When specializing $A = \mathbb{Z}$, we have that $\mathcal{D}(R)/p\mathcal{D}(R) = \mathcal{D}(R/pR)$.

Remark 4.2. Let R be a noetherian commutative ring of characteristic p . Assume that R is a finitely generated R^p -module. Then by [Yek92, 1.4.9] and [SVdB97, 2.5.1] we have

$$\mathcal{D}(R) = \mathcal{D}(R, \mathbb{Z}/p\mathbb{Z}) = \bigcup_e \text{Hom}_{R^{p^e}}(R, R).$$

Proposition 4.3. *Let R be a regular ring of characteristic p such that R is a finitely generated R^p -module. Then each F_R -module is a $\mathcal{D}(R)$ -module.*

Proof. Let \mathcal{M} be an F_R -module with the structure isomorphism $\theta : \mathcal{M} \rightarrow F_R(\mathcal{M})$. By Remark 4.2, it suffices to specify the action of each element $\delta \in \text{Hom}_{R^{p^e}}(R, R)$ on \mathcal{M} . Let θ_e denote the composition $F^{e-1}(\theta) \circ \dots \circ \theta : \mathcal{M} \rightarrow F^e(\mathcal{M})$. Clearly δ acts on $F^e(\mathcal{M}) = F_*^e R \otimes_R \mathcal{M}$ via $\delta \otimes \text{id}_{\mathcal{M}}$. We let δ act on \mathcal{M} via $\theta_e^{-1} \circ (\delta \otimes \text{id}_{\mathcal{M}}) \circ \theta_e$.

Since $\delta \in \text{Hom}_{R^{p^e}}(R, R) \subseteq \text{Hom}_{R^{p^{e'}}}(R, R)$ for $e' > e$, we need to check that the action of δ is independent of the choice of e , *i.e.* we need to check that

$$(1) \quad \theta_e^{-1} \circ (\delta \otimes \text{id}_{\mathcal{M}}) \circ \theta_e = \theta_{e'}^{-1} \circ (\delta \otimes \text{id}_{\mathcal{M}}) \circ \theta_{e'}.$$

Set $\theta_{e'-e}^e = F^e(\theta_{e'-e}) = \text{id}_{F_*^e R} \otimes \theta_{e'-e}$. Then $\theta_{e'} = \theta_{e'-e}^e \circ \theta_e$. To check (1) is equivalent to checking

$$\begin{aligned} \delta \otimes \text{id}_{\mathcal{M}} &= (\theta_{e'-e}^e)^{-1} \circ (\delta \otimes \text{id}_{\mathcal{M}}) \circ \theta_{e'-e}^e \\ &= (\text{id}_{F_*^e R} \otimes \theta_{e'-e})^{-1} \circ (\delta \otimes \text{id}_{\mathcal{M}}) \circ (\text{id}_{F_*^e R} \otimes \theta_{e'-e}) \end{aligned}$$

which is clear. \square

As an application of \mathcal{D} -module, now we turn our attention to answering the question raised in Remark 3.10.

Let $R = \mathbb{F}_p[x_1, \dots, x_n]$ and let \mathcal{D} denote the ring of differential operators over R . We will use $\partial_j^{[t]}$ to denote $\frac{1}{t!} \frac{\partial^t}{\partial x_j^t}$.

Definition 4.4. The r -th Euler operator, denoted by E_r , is defined as

$$E_r := \sum_{i_1+i_2+\dots+i_n=r, i_1 \geq 0, \dots, i_n \geq 0} x_1^{i_1} \dots x_n^{i_n} \partial_1^{[i_1]} \dots \partial_n^{[i_n]}.$$

In particular E_1 is the usual Euler operator $\sum_{i=1}^n x_i \partial_i$.

A graded \mathcal{D} -module M is called *Eulerian*, if each homogeneous element $z \in M$ satisfies

$$(2) \quad E_r \cdot z = \binom{\deg(z)}{r} \cdot z$$

for every $r \geq 1$.

Remark 4.5. An F_R -module \mathcal{M} with structure isomorphism $\theta : \mathcal{M} \rightarrow F_R(\mathcal{M})$ is called a *graded F_R -module* if \mathcal{M} is a graded R -module and θ is degree-preserving. It turns out that each graded F_R -module is an Eulerian graded \mathcal{D} -module [MZ14, Theorem 4.4].

Exercise 4.6. Prove that, if M is a graded Eulerian \mathcal{D} -module, so are \mathcal{D} -submodules and \mathcal{D} -quotients of M .

Exercise 4.7. Prove that M_f is an Eulerian graded \mathcal{D} -module for an Eulerian graded \mathcal{D} -module M and a graded element $f \in R$ and \cdot . (Hint: $\partial_i^{[j]}$ is p^e -linear for $p^e > j$ and hence $\partial_i^{[j]}(\frac{z}{f^{p^e}}) = \frac{1}{f^{p^e}} \partial_i^{[j]}(z)$ for each $z \in M$.)

Combining Exercises 4.6 and 4.7, we have

Proposition 4.8. Let $\mathfrak{a}_1, \dots, \mathfrak{a}_s$ be graded ideals of R . Then each local cohomology module $H_{\mathfrak{m}}^i H_{\mathfrak{a}_1}^{j_1} \dots H_{\mathfrak{a}_s}^{j_s}(R)$ is an Eulerian graded \mathcal{D} -module.

For each graded R -module M and integer ℓ , we will use $M(\ell)$ to denote the graded R -module whose degree- j piece is $M_{j+\ell}$ for each integer j .

Exercise 4.9. Let M be an Eulerian graded \mathcal{D} -module. Prove that $M(\ell)$ is Eulerian if and only if $\ell = 0$.

Remark 4.10. The ring of differential operators \mathcal{D} has a natural grading $\deg(x_i) = 1$ and $\deg(\partial_i^{[j]}) = -j$. Under this grading, the map $\frac{\mathcal{D}}{\mathcal{D}_{\mathfrak{m}}} \rightarrow H_{\mathfrak{m}}^n(R)(-n)$ defined by

$$\partial_1^{[i_1]} \dots \partial_n^{[i_n]} \mapsto (-1)^{i_1 + \dots + i_n} x_1^{-i_1 - 1} \dots x_n^{-i_n - 1}$$

is a degree-preserving isomorphism.

Theorem 4.11. *If M is an Eulerian graded \mathcal{D} -module and $\text{Supp}_R(M) = \{\mathfrak{m}\}$, then M is isomorphic (as graded R -modules) to a direct sum of copies of $H_{\mathfrak{m}}^n(R)$.*

Proof. Let $\text{Soc}(M)$ denote the socle of M , i.e. the sub- R -module of M that is annihilated by \mathfrak{m} . It is straightforward to check that $\text{Soc}(M)$ is generated by graded elements and a minimal set of graded generators of $\text{Soc}(M)$ is a graded k -basis for $\text{Soc}(M)$.

Let $\{\beta_j\}$ be a graded k -basis of $\text{Soc}(M)$ with $\deg(\beta_j) = -n_j$. There is a degree-preserving homomorphism of \mathcal{D} -modules $\bigoplus_j \frac{\mathcal{D}}{\mathcal{D}_{\mathfrak{m}}}(n_j) \rightarrow M$ which sends 1 of the j -th copy to β_j . This map is an isomorphism because it induces an isomorphism on socles. By Remark 4.10, we have $\bigoplus_j \frac{\mathcal{D}}{\mathcal{D}_{\mathfrak{m}}}(n_j) \cong \bigoplus_j H_{\mathfrak{m}}^n(R)(-n + n_j)$. Hence $M \cong \bigoplus_j H_{\mathfrak{m}}^n(R)(-n + n_j)$. Since M is Eulerian, so is each $H_{\mathfrak{m}}^n(R)(-n + n_j)$ which implies that $n_j = n$ for each j by Exercise 4.9. This finishes the proof. \square

5. INTERACTIONS BETWEEN F -MODULES AND \mathcal{D} -MODULES

We have seen in Proposition 4.3 that each F_R -module is a $\mathcal{D}(R)$ -module. It turns out that the theories of F -modules and \mathcal{D} -modules are intertwined and their interactions have had many applications. We begin with the following result due to Lyubeznik [Lyu97, 5.7].

Theorem 5.1. *Let R be a regular ring of characteristic p that is a finite R^p -module. Then each F_R -finite module has finite length in the category of $\mathcal{D}(R)$ -modules.*

We will see an application of this theorem in a moment.

Exercise 5.2. Let $\phi : R \rightarrow S$ be a ring homomorphism between commutative rings such that locally S is a finitely generated free R -module. Then the category of R -modules is equivalent to the one of $\text{End}_R(S)$ -modules. The functors are $S \otimes_R -$ and $\text{Hom}_R(S, R) \otimes_{\text{End}_R(S)} -$.

Remark 5.3. We'd like to apply the equivalence in Exercise 5.2 to Frobenius endomorphism.

Let R be a regular ring of characteristic p that is a finite R^p -module. Consider the e -th Frobenius $R \rightarrow F_*^e R$ (here we denote the target ring by $F_*^e R$ assuming no confusion will arise). Clearly $\text{End}_R(F_*^e R) \cong \text{End}_{R^{p^e}}(R)$ (keep in mind that this is part of the ring $\mathcal{D}(R)$ according to Remark 4.2). Hence the category of $\text{End}_{R^{p^e}}(R)$ -modules is equivalence to the one of R -modules.

One immediate consequence is that the categories of $\text{End}_{R^{p^e}}(R)$ -modules are equivalent for all $e \geq 1$. The functor that gives the equivalence between the category of $\text{End}_{R^{p^e}}(R)$ -modules and the one of $\text{End}_{R^{p^{e+e'}}}(R)$ -modules is $F_R^{e'}(-) = F_*^{e'} R \otimes_R -$. More specifically, let M be a $\text{End}_{R^{p^e}}(R)$ -module. The $\text{End}_{R^{p^{e+e'}}}(R)$ -module structure on $F_R^{e'}(M)$ can be seen as follows. Since the category of $\text{End}_{R^{p^e}}(R)$ -modules is equivalent to the one of R -modules, we know that $M \cong F_*^e R \otimes_R N$ for some R -module N .

Then $F_R^{e'} \otimes_R M \cong F_*^{e+e'} R \otimes_R N$. And hence $\text{End}_{R^{pe+e'}}(R)$ acts on $F_R^{e'}(M)$ via its action on $F_*^{e+e'} R \otimes_R N$ which is given by $\alpha \otimes id_N$ for each $\alpha \in \text{End}_{R^{pe+e'}}(R)$.

By our discussion so far, we can see that, when R is a regular ring of characteristic p that is a finite R^p -module, the functor $F_R^e(-)$ is an equivalence of the category of $\mathcal{D}(R)$ -modules to itself.

Theorem 5.4. *Let R be a regular ring of characteristic p that is a finite R^p -module. Assume that \mathcal{M} is an F_R -finite module and M is a root of \mathcal{M} . Then the image of M in \mathcal{M} generates \mathcal{M} as a $\mathcal{D}(R)$ -module.*

Proof. Since M is a root of \mathcal{M} , we know that $\beta : M \rightarrow F_R(M)$ is injective and hence the direct limit (which is \mathcal{M}) of $M \rightarrow F_R(M) \rightarrow F_R^2(M) \rightarrow \dots$ is a direct union. Thus, we may assume that $M \subset F_R(M)$ and $\mathcal{M} = \bigcup_e F_R^e(M)$.

Let \mathcal{M} be the $\mathcal{D}(R)$ -submodule of \mathcal{M} that is generated by M . Since $M \subset F_R(M)$, we have $\mathcal{M} \subset F_R(\mathcal{M})$. If \mathcal{M} were strictly contained in $F_R(\mathcal{M})$ (which is also a $\mathcal{D}(R)$ -submodule of \mathcal{M}), then we would have a strictly increasing chain of $\mathcal{D}(R)$ -submodules of \mathcal{M} :

$$\mathcal{M} \subsetneq F_R(\mathcal{M}) \subsetneq F_R^2(\mathcal{M}) \subsetneq \dots$$

which is a contradiction to Theorem 5.1. Therefore $\mathcal{M} = F_R(\mathcal{M})$. Consequently $\mathcal{M} = \bigcup_e F_R^e(M) \subseteq \bigcup_e F_R^e(\mathcal{M}) = \mathcal{M}$. This finishes the proof. \square

Corollary 5.5. *Let R be a regular ring of characteristic p that is a finite R^p -module. Assume that \mathcal{M} is an F_R -finite module and $\beta : M \rightarrow F_R(M)$ is a generating morphism of \mathcal{M} . Then the image of M in \mathcal{M} generates \mathcal{M} as a $\mathcal{D}(R)$ -module.*

As an application of the interactions between F -modules and \mathcal{D} -modules, we prove the following result [BBL⁺14, 3.1(1)].

Theorem 5.6. *Let $R = \mathbb{Z}[x_1, \dots, x_n]$ and \mathfrak{a} be an ideal of R . If a prime number p is a nonzerodivisor on $\text{Ext}_R^i(R/\mathfrak{a}, R)$, then p is also a nonzerodivisor on $H_{\mathfrak{a}}^i(R)$.*

Proof. Let p be a prime integer. The exact sequence $0 \rightarrow R \xrightarrow{p} R \rightarrow R/pR \rightarrow 0$ induces an exact sequence of Ext and an exact sequence of local cohomology modules; they fit into a commutative diagram:

$$\begin{array}{ccccccc} \text{Ext}_R^{i-1}(R/\mathfrak{a}, R) & \xrightarrow{\psi} & \text{Ext}_R^{i-1}(R/\mathfrak{a}, R/pR) & \longrightarrow & \text{Ext}_R^i(R/\mathfrak{a}, R) & \xrightarrow{p} & \text{Ext}_R^i(R/\mathfrak{a}, R) \\ \downarrow \beta & & \downarrow \gamma & & \downarrow & & \\ H_{\mathfrak{a}}^{i-1}(R) & \xrightarrow{\varphi} & H_{\mathfrak{a}}^{i-1}(R/pR) & \xrightarrow{\delta} & H_{\mathfrak{a}}^i(R) & \xrightarrow{p} & H_{\mathfrak{a}}^i(R) \end{array}$$

The bottom row is a complex of $\mathcal{D}(R)$ -modules; in particular, $\varphi(H_{\mathfrak{a}}^{i-1}(R))$ is a $\mathcal{D}(R)$ -submodule of $H_{\mathfrak{a}}^{i-1}(R/pR)$. As $\varphi(H_{\mathfrak{a}}^{i-1}(R))$ is annihilated by p , it has a natural structure as a module over the ring $\mathcal{D}(R)/p\mathcal{D}(R)$, which equals $\mathcal{D}(R/pR)$. Similarly,

$$(3) \quad H_{\mathfrak{a}}^{i-1}(R/pR) \xrightarrow{\delta} \text{Im}(\delta)$$

is a map of $\mathcal{D}(R/pR)$ -modules.

Suppose p is a nonzerodivisor on $\text{Ext}_R^i(R/\mathfrak{a}, R)$. Then the map ψ is surjective; we need to prove that p is a nonzerodivisor on $H_{\mathfrak{a}}^i(R)$, equivalently, that φ is surjective.

By Corollary 5.5, the image M of γ generates $H_{\mathfrak{a}}^{i-1}(R/pR)$ as a $\mathcal{D}(R/pR)$ -module. As ψ is surjective, M is also the image of $\gamma \circ \psi = \varphi \circ \beta$. It follows that

$$M \subseteq \varphi(H_{\mathfrak{a}}^{i-1}(R)).$$

But $\varphi(H_{\mathfrak{a}}^{i-1}(R))$ is a $\mathcal{D}(R/pR)$ -submodule of $H_{\mathfrak{a}}^{i-1}(R/pR)$ that contains M . Hence

$$\varphi(H_{\mathfrak{a}}^{i-1}(R)) = H_{\mathfrak{a}}^{i-1}(R/pR),$$

i.e., φ is surjective, as desired. \square

Corollary 5.7. *Let $R = \mathbb{Z}[x_1, \dots, x_n]$ and \mathfrak{a} be an ideal of R . Then there are only finitely many prime numbers p such that $H_{\mathfrak{a}}^j(R)$ contains a nonzero p -torsion element for all j .*

We have seen in Theorem 3.14 that F_R -finite modules have finite length in the category of F_R -module. On the other hand, [Lyu97, 5.7] says that an F_R -finite module with the \mathcal{D} -module structure given in Proposition 4.3 also has finite length in the category of \mathcal{D} -modules. So, a natural question is how does one compare these two lengths. It turns out that they are the same when the underlying field is algebraically closed.

Theorem 5.8 (Theorem 1.1 in [Bli03]). *Let R be a noetherian regular ring that is essentially of finite type over an algebraically closed field of characteristic p . Then the length of an F_R -finite module in the category of F -modules is the same as its length in the category of \mathcal{D} -modules.*

Question 5.9. Let \mathfrak{a} be an ideal of R . Can one compute the length of $H_{\mathfrak{a}}^i(R)$ as a \mathcal{D} -module?

At the moment, very little is known about Question 5.9; even when \mathfrak{a} is a principal ideal and $i = 1$. We will come back to this question when we discuss non-regular rings in characteristic p .

6. RING-THEORETIC PROPERTIES CHARACTERIZED BY FROBENIUS

Theorem 0.5 exemplifies that the Frobenius endomorphism encodes properties of the ring itself.

Definition 6.1. R is called *F -pure* if $M \rightarrow F_*R \otimes_R M$ is injective for all R -modules M .

When (R, \mathfrak{m}) is local, R is called *F -injective* if the Frobenius action $H_{\mathfrak{m}}^i(R) \xrightarrow{F} H_{\mathfrak{m}}^i(R)$ is injective all i . More generally when R may not be local, R is called *F -injective* if $R_{\mathfrak{p}}$ is so for each prime ideal \mathfrak{p} .

Exercise 6.2. Let (R, \mathfrak{m}) be a noetherian local ring of characteristic p . Let $E(R/\mathfrak{m})$ denote the injective hull of R/\mathfrak{m} . Prove that R is F -pure if and only if $E(R/\mathfrak{m}) \rightarrow F_*R \otimes_R E(R/\mathfrak{m})$ is injective.

Exercise 6.3. Let (R, \mathfrak{m}) be a noetherian local ring of characteristic p . Assume that R is a finite R^p -module. Prove that the following are equivalent

- (1) R is F -pure;
- (2) for each $e \geq 1$, there is an R^{p^e} -module homomorphism $\varphi : R \rightarrow R^{p^e}$ such that the composition $R^{p^e} \hookrightarrow R \xrightarrow{\varphi} R^{p^e}$ is identity on R^{p^e} ;
- (3) for some $e \geq 1$, there is an R^{p^e} -module homomorphism $\varphi : R \rightarrow R^{p^e}$ such that the composition $R^{p^e} \hookrightarrow R \xrightarrow{\varphi} R^{p^e}$ is identity on R^{p^e} .

Example 6.4. Let $R = k[[x_1, \dots, x_d]]$ and $f \in \mathfrak{m} = (x_1, \dots, x_d)$ be a formal power series. Since R/fR is a hypersurface and hence Gorenstein, $E((R/fR)/\mathfrak{m}) = H_{\mathfrak{m}}^{d-1}(R/fR)$. So, R/fR is F -pure if and only if the Frobenius action $F : H_{\mathfrak{m}}^{d-1}(R/fR) \rightarrow H_{\mathfrak{m}}^{d-1}(R/fR)$ is injective. Consider the following commutative diagram induced by $0 \rightarrow R \xrightarrow{f} R \rightarrow R/fR \rightarrow 0$:

$$(4) \quad \begin{array}{ccccccc} 0 & \longrightarrow & H_{\mathfrak{m}}^{d-1}(R/fR) & \longrightarrow & H_{\mathfrak{m}}^d(R) & \xrightarrow{f} & H_{\mathfrak{m}}^d(R) \longrightarrow 0 \\ & & \downarrow F & & \downarrow f^{p-1}F & & \downarrow F \\ 0 & \longrightarrow & H_{\mathfrak{m}}^{d-1}(R/fR) & \longrightarrow & H_{\mathfrak{m}}^d(R) & \xrightarrow{f} & H_{\mathfrak{m}}^d(R) \longrightarrow 0 \end{array}$$

It is easy to see from this diagram that $F : H_{\mathfrak{m}}^{d-1}(R/fR) \rightarrow H_{\mathfrak{m}}^{d-1}(R/fR)$ is injective if the map in the middle $f^{p-1}F$ is injective. From Example 0.3, we can see that this is the case if and only if $f^{p-1} \notin \mathfrak{m}^{[p]}$.

More generally, the quotient ring R/I is F -pure if and only if $(I^{[p]} : I) \not\subseteq \mathfrak{m}^{[p]}$; this is called Fedder's Criterion.

Exercise 6.5. Let $R = \frac{\mathbb{F}_p[[x,y,z]]}{(x^3+y^3+z^3)}$ and consider the element $[\frac{z^2}{xy}] \in H_{(x,y,z)}^2(R)$. Find all prime numbers p such that the image of $[\frac{z^2}{xy}]$ under the Frobenius action on $H_{(x,y,z)}^2(R)$ is 0.

Question 6.6. Assume that f is a polynomial of degree d in d variables x_1, \dots, x_d with integer coefficients and that its Jacobian ideal is primary to (x_1, \dots, x_d) . Let f_p denote the image of f under $R := \mathbb{Z}[[x_1, \dots, x_d]] \rightarrow R_p := \mathbb{F}_p[[x_1, \dots, x_d]]$. Are there infinitely many prime numbers p such that $R_p/f_p R_p$ is F -pure?

Remark 6.7. When f is a degree 3 homogeneous polynomial that defines an elliptic curve, then by [Sil92, 5.11 on page 145] there are infinitely many prime numbers p such that f_p defines an ordinary elliptic curve and hence $R_p/f_p R_p$ is F -pure.

Exercise 6.8. Assume that (R, \mathfrak{m}) is an F -injective local ring. Prove that R is reduced.

Question 6.9. Let f be a regular element of a local ring (R, \mathfrak{m}) . Assume that R/fR is F -injective. Is it true that R is F -injective?

Exercise 6.10. Let (R, \mathfrak{m}) be Cohen-Macaulay and $f \in R$ be a regular element. Assume that R/fR is F -injective. Prove that R is also F -injective.

Question 6.9 is still open in its full generality. For recent developments, see [HMS14].

Definition 6.11. Let R be a noetherian commutative ring of characteristic p . We say that R is strongly F -regular if for every non-zero element $r \in R$ there exists $e \in \mathbb{N}$ and $\phi \in \text{Hom}_{R^{p^e}}(R, R^{p^e})$ such that $\phi(r) = 1$.

Exercise 6.12. Prove that, if R is a strongly F -regular local ring, then R is a domain.

Exercise 6.13. Let $R \subset S$ be a split ring extension (*i.e.* there is an R -module homomorphism $S \rightarrow R$ such that $R \rightarrow S \rightarrow R$ is the identity on R). Assume that S is strongly F -regular. Prove that R is also strongly F -regular.

Question 6.14. Let R be a noetherian commutative ring of characteristic p . An ideal I of R is called *Frobenius closed* if $r^p \in I^{[p]}$ implies $r \in I$ for all elements $r \in R$. Assume now that R is local and consider the following two conditions

- (1) R is F -injective;
- (2) each parameter ideal I is Frobenius closed.

Is it true that (1) and (2) are equivalent?

Question 6.15. Does the conclusion of Exercise 1.1 still hold when assuming R is strongly F -regular instead of regular?

It turns out that one can use properties of \mathcal{D} -modules to characterize strong F -regularity.

Theorem 6.16 (Theorem 2.2(4) in [Smi95]). *Let (R, \mathfrak{m}) be a noetherian commutative local ring of characteristic p . Assume that R is a finite R^p -module. Then R is strongly F -regular if and only if R is F -pure and a simple \mathcal{D} -module.*

Proof. First assume that R is strongly F -regular. Then it is clear that R is F -pure. We will show that $\mathcal{D}(R)r = R$ for each nonzero element $r \in R$. Since R is strongly F -regular, there is an integer $e \geq 1$ and a $\varphi \in \text{Hom}_{R^{p^e}}(R, R^{p^e})$ such that $\varphi(r) = 1$. The composition of φ with $R^{p^e} \hookrightarrow R$, still denoted by φ , is clearly an element of $\text{End}_{R^{p^e}}(R) \subset \mathcal{D}(R)$. Since $\varphi(r) = 1$, we have $\mathcal{D}(R)r = R$.

Conversely, assume that R is F -pure and a simple \mathcal{D} -module. Let r be a nonzero element of R . Since R is $\mathcal{D}(R)$ -simple, there is $\delta \in \mathcal{D}(R)$ such that $\delta(r) = 1$. By Remark 4.2, δ is R^{p^e} -linear for some $e \geq 1$. By Exercise 6.3, there is a $\varphi \in \text{Hom}_{R^{p^e}}(R, R^{p^e})$ that splits the inclusion $R^{p^e} \hookrightarrow R$. Now $\varphi \circ \delta : R \rightarrow R^{p^e}$ is R^{p^e} -linear and $\varphi \circ \delta(r) = 1$. \square

Question 6.17. Let R be an integral domain of characteristic p that is a finite R^p -module. Assume that every module-finite ring extension $R \hookrightarrow S$ splits. Is it true that R is $\mathcal{D}(R)$ -simple?

The following result due to Blickle [Bli04, 4.10] sheds some light on Question 5.9.

Theorem 6.18. *Let (R, \mathfrak{m}) be an F -finite regular local ring and $f \in \mathfrak{m}$ be an element of R . Assume that R/fR is F -injective (or equivalently, F -pure). Then $H_{(f)}^1(R)$ is a simple $\mathcal{D}(R)$ -module if and only if R/fR is strongly F -regular.*

REFERENCES

- [BBL⁺14] B. BHATT, M. BLICKLE, G. LYUBEZNIK, A. K. SINGH, AND W. ZHANG: *Local cohomology modules of a smooth \mathbb{Z} -algebra have finitely many associated primes*, Invent. Math. **197** (2014), no. 3, 509–519. 3251828
- [Bli03] M. BLICKLE: *The D -module structure of $R[F]$ -modules*, Trans. Amer. Math. Soc. **355** (2003), no. 4, 1647–1668. 1946409 (2004m:13016)
- [Bli04] M. BLICKLE: *The intersection homology D -module in finite characteristic*, Math. Ann. **328** (2004), no. 3, 425–450. 2036330 (2005a:14005)
- [Har68] R. HARTSHORNE: *Cohomological dimension of algebraic varieties*, Ann. of Math. (2) **88** (1968), 403–450. 0232780 (38 #1103)
- [HS77] R. HARTSHORNE AND R. SPEISER: *Local cohomological dimension in characteristic p* , Ann. of Math. (2) **105** (1977), no. 1, 45–79. MR0441962 (56 #353)
- [HE71] M. HOCHSTER AND J. A. EAGON: *Cohen-Macaulay rings, invariant theory, and the generic perfection of determinantal loci*, Amer. J. Math. **93** (1971), 1020–1058. 0302643 (46 #1787)
- [HMS14] J. HORIUCHI, L. E. MILLER, AND K. SHIMOMOTO: *Deformation of F -injectivity and local cohomology*, Indiana Univ. Math. J. **63** (2014), no. 4, 1139–1157. 3263925
- [Hun92] C. HUNEKE: *Problems on local cohomology*, Free resolutions in commutative algebra and algebraic geometry (Sundance, UT, 1990), Res. Notes Math., vol. 2, Jones and Bartlett, Boston, MA, 1992, pp. 93–108. 1165320 (93f:13010)

- [HK91] C. HUNEKE AND J. KOH: *Cofiniteness and vanishing of local cohomology modules*, Math. Proc. Cambridge Philos. Soc. **110** (1991), no. 3, 421–429. 1120477 (92g:13021)
- [HS93] C. L. HUNEKE AND R. Y. SHARP: *Bass numbers of local cohomology modules*, Trans. Amer. Math. Soc. **339** (1993), no. 2, 765–779. 1124167 (93m:13008)
- [Kun69] E. KUNZ: *Characterizations of regular local rings for characteristic p* , Amer. J. Math. **91** (1969), 772–784. 0252389 (40 #5609)
- [Lyu97] G. LYUBEZNIK: *F -modules: applications to local cohomology and D -modules in characteristic $p > 0$* , J. Reine Angew. Math. **491** (1997), 65–130. MR1476089 (99c:13005)
- [Lyu06] G. LYUBEZNIK: *On the vanishing of local cohomology in characteristic $p > 0$* , Compos. Math. **142** (2006), no. 1, 207–221. 2197409 (2007b:13029)
- [MZ14] L. MA AND W. ZHANG: *Eulerian graded \mathcal{D} -modules*, Math. Res. Lett. **21** (2014), no. 1, 149–167. 3247047
- [PS73] C. PESKINE AND L. SZPIRO: *Dimension projective finie et cohomologie locale. Applications à la démonstration de conjectures de M. Auslander, H. Bass et A. Grothendieck*, Inst. Hautes Études Sci. Publ. Math. (1973), no. 42, 47–119. 0374130 (51 #10330)
- [Sil92] J. H. SILVERMAN: *The arithmetic of elliptic curves*, Graduate Texts in Mathematics, vol. 106, Springer-Verlag, New York, 1992, Corrected reprint of the 1986 original. 1329092 (95m:11054)
- [Sin00] A. K. SINGH: *p -torsion elements in local cohomology modules*, Math. Res. Lett. **7** (2000), no. 2-3, 165–176. 1764314 (2001g:13039)
- [Smi95] K. E. SMITH: *The D -module structure of F -split rings*, Math. Res. Lett. **2** (1995), no. 4, 377–386. 1355702 (96j:13024)
- [SVdB97] K. E. SMITH AND M. VAN DEN BERGH: *Simplicity of rings of differential operators in prime characteristic*, Proc. London Math. Soc. (3) **75** (1997), no. 1, 32–62. 1444312 (98d:16039)
- [Yek92] A. YEKUTIELI: *An explicit construction of the Grothendieck residue complex*, Astérisque (1992), no. 208, 127, With an appendix by Pramathanath Sastry. 1213064 (94e:14026)